

Apparent horizons, black hole entropy and loop quantum gravity

Viqar Husain*

*Department of Physics and Astronomy,
University of British Columbia,
6224 Agricultural Road
Vancouver, BC V6T1Z1, Canada*

Abstract

In recent work on black hole entropy in non-perturbative quantum gravity, an action for the black hole sector of the phase space is introduced and (partially) quantized. We give a number of observations on this and related works. In particular we show that (i) the entropy calculation applies without change to generally covariant theories having no black hole solutions, (ii) the phase space constraint used to select the black hole sector is not the apparent horizon equation, which is the natural phase space constraint separating trapped and untrapped regions on an initial data surface, and (iii) there appears to be at least one other phase space constraint which leads to the conclusion that the entropy associated with a bounding two-dimensional surface is proportional to its area.

PACS numbers:

Typeset using REVTeX

*Email address: husain@physics.ubc.ca

Among the latest developments in the non-perturbative approach to quantum gravity utilizing the triad-connection canonical variables [1–6], is a set of ideas that help to answer the question: what are the microscopic degrees of freedom of a black-hole? Very briefly, the answer in this approach is that black hole microstates are certain “surface” degrees of freedom residing on the horizon of a black hole. Counting of these degrees of freedom yields an entropy S_{BH} proportional to the area A of the black hole event horizon for large black holes, in accord with the Beckenstein-Hawking formula; the proportionality constant in the proposal contains a parameter, which it is argued may be fixed to give $S_{BH} = A/4$ in Planck units.

The initial concrete realization of the idea that surface degrees of freedom residing on the event horizon may be identified with black hole microstates was given by Carlip [7] for the three-dimensional black hole.¹ In four dimensions, the stage for calculating black hole entropy as a counting of surface degrees of freedom was set by work of Smolin [8] on the Beckenstein bound, and of Rovelli [9] and Krasnov [10] on counting the degeneracy of spin network states associated with fixed area.

The four-dimensional black hole entropy calculation due to Ashtekar, Baez, Corichi and Krasnov [11] in the loop quantum gravity approach is similar in spirit to these earlier works. It contains a concrete proposal for an action for the “black hole sector” of general relativity. This is used as the starting point for canonical quantization.

In this note we give an outline review of the background and main ingredients of this proposal, and then give a number of observations on them. These are: (I) a nearly identical analysis may be carried out in theories without “dynamics,” which have no black hole solutions, with the resulting entropy having a different interpretation, namely, that an entropy (proportional to area) may be associated to any surface, (II) the phase space condition used in selecting the “black hole sector” of the gravitational phase space in [11] is not the apparent horizon equation, which is the phase space condition giving the boundary separating trapped and untrapped regions on a spatial slice, and (III) other boundary conditions may also give a surface entropy proportional to the surface area.

The non-perturbative quantum gravity approach starts with a classical first order “self-dual” action in which the gravitational degrees of freedom are the $SL(2, \mathbb{C})$ vierbein $e_a^{AA'}$, and self-dual part of the $SL(2, \mathbb{C})$ connection A_{aB}^A . This connection has only chiral 2-spinor components, and is therefore complex [12]. Boundary terms may be added to the bulk action as required by the physical context to guarantee a well-defined variational principle.

The main ingredients in the black hole entropy calculation are as follows; (the reader is referred the original papers [11], and references therein for further details).

(i) Attention is restricted to non-static and asymptotically flat black hole spacetimes M with an inner boundary (the event horizon H). A specific phase space condition is specified on this boundary, and is incorporated into the following proposed action for black hole spacetimes:

$$S[e, A] = \frac{1}{8\pi G_N} \int_M \text{Tr} [e \wedge e \wedge F(A)] + \frac{1}{8\pi G_N} \frac{A_S}{4\pi} \int_H \text{Tr} \left[A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right], \quad (1)$$

¹For a recent reanalysis of this work see [28].

where A_S is the area of the 2-surface boundary S where spatial slices Σ intersect the horizon. The boundary condition which makes this action functionally differentiable is ²

$$\underline{F}_{ab} = -\frac{2\pi}{A_S} \underline{\sigma}_{ab}, \quad (2)$$

where $\sigma_{ab} = (e \wedge e)_{ab}$ is the dual of the momentum E_i^a conjugate to A_a^i , and the underline denotes restrictions of the fields to S ; $i = 1, 2, 3$ is the $SU(2)$ index). Since the surface term in this action contains time derivatives, there is an additional surface contribution to the phase space Poisson brackets, and hence dynamical surface degrees of freedom.

(ii) This action leads to the complex phase space configuration variable $A_a^i = \Gamma_a^i - iK_a^i$, where Γ_a^i is the spatial spin connection and K_a^i is the extrinsic curvature. At this stage a change to real variables is made [13–15]: this variable is replaced with the real phase space variable ${}^\gamma A_a := \Gamma_a^i - \gamma K_a^i$, where γ is a real parameter (the Barbero-Immirzi parameter [15]). Its conjugate momentum variable is ${}^\gamma \tilde{E}^a = (1/\gamma)\tilde{E}^a$. The horizon boundary condition on the two-surface S in Σ is now

$${}^\gamma \underline{F}_{ab} = -\frac{2\pi\gamma}{A_S} {}^\gamma \underline{\sigma}_{ab}, \quad (3)$$

where both sides of the equation are real. This (γ dependent) equation is a part of the conditions used to identify the black hole sector of the phase space, and is the one imposed quantum mechanically. (The remaining conditions in [11] are not directly relevant for the purposes of this paper.)

(iii) The kinematical constraints arising from the above action, which generate the $SU(2)$ Gauss law and spatial diffeomorphisms, can be quantized; the kinematical quantum states $\psi[A]$ are spin network states [16,17]. These are basically the generalization of Wilson lines to graphs, with edges labelled by $SU(2)$ representations, and group index contractions at the vertices via generalizations of 6-j and 9-j etc. symbols. In the black hole context, the full Hilbert space is a direct product of “volume” and “surface” states: $\psi[A] = \psi_V[A] \otimes \psi_S[A]$.

(iv) The horizon boundary condition in (2) is imposed as the following quantum condition to select out kinematical black hole states:

$$\left(1 \otimes \frac{2\pi\gamma}{A_S} \hat{\underline{F}}_{ab} \cdot r + \hat{\underline{\sigma}}_{ab} \cdot r \otimes 1\right) \Psi_V \otimes \Psi_S = 0 \quad (4)$$

where r is a fixed internal vector. The first and second terms act entirely on surface and volume states respectively. The volume states are spin network states, and the surface states are states of Chern-Simons theory with sources provided by spin network edges intersecting the surface [11].

(v) Spin network states are also eigenstates of the area operator [18,19]. Therefore, the relevant solutions of (4) are those spin network states which are compatible with the fixed area A_S in (4). The entropy is calculated by counting the number of solutions of (4) with this added restriction. This counting gives entropy proportional to A_S .

We now give three observations and related results on this black hole entropy calculation:

²For details of the origin of this and other conditions see [11].

(I) These steps are entirely kinematical in the sense that the Hamiltonian constraint, or its consequences, do not enter any of the calculations: the space of states used are solutions to the Gauss and spatial diffeomorphism constraints, and the phase space condition (2) is also independent of the bulk dynamics (as any phase space condition other than the Hamiltonian constraint must be). Thus, it is clear that the main steps outlined above may be carried out for *any* theory which has the same kinematics, whether or not it has a Hamiltonian, or Hamiltonian constraint. What must be different however, is the interpretation to be attached to the calculation if the theory is not the black hole sector of general relativity (or any other gravitational theory with black hole solutions for that matter).

This may be illustrated with a concrete example: A generally covariant theory with the same kinematics as general relativity, but having no black hole solutions, is obtained from an action identical to the bulk action (1) above, but with $SU(2)$ rather than $SL(2, \mathbb{C})$ as the gauge group [20]. The (real) covariant dynamical fields are e_a^i and A_a^i , where $i = 1, 2, 3$ is the $SU(2)$ index. The Chern-Simons surface term may be added to the action to make the analogy complete. Thus, consider the action

$$S[e, A] = \frac{1}{8\pi\lambda} \int_M (e^i \wedge e^j \wedge F^k(A) \epsilon_{ijk}) + \frac{1}{8\pi\lambda} \frac{A_S}{4\pi} \int_{\partial M} \left(A^i \wedge dA^i + \frac{2}{3} A^i \wedge A^j \wedge A^k \epsilon_{ijk} \right) \quad (5)$$

on a manifold M with boundary ∂M . The constant A_S is now taken to be the area of the 2-surface S obtained by the intersection of ∂M with a spatial surface Σ in M , (which is an embedded surface in M on which the induced metric is not degenerate). The action (5) has a coupling constant λ . Its dimension is fixed by assuming that e_a^i is dimensionless and A_a^i has dimension $(\text{length})^{-1}$, so $[\lambda] = \text{time/mass}$ (giving S dimensions of action). Note that there is no speed of light c in the theory because the “spacetime” metric $g_{ab} = e_a^i e_b^j \delta_{ij}$ is degenerate with signature $(0+++)$; e_a^i is a dreibein in four dimensions. Therefore, although the quantum theory has a fundamental length $l_F = \sqrt{\hbar\lambda}$, it does not have a fundamental mass or time.

No component of the boundary ∂M is a horizon. Indeed, no spacetime horizon can even be defined because the four-metric is degenerate. Nevertheless, the boundary condition on ∂M required by functional differentiability of the action is identical to (2) above, as may be directly verified.

We are free to define a boundary with more than one component. Therefore consider spatial slices Σ on which there is an inner boundary S of area A_S , and an outer asymptotic region. Hamiltonian decomposition of this generally covariant $SU(2)$ gauge theory reveals, identically to the gravitational case, that there are volume *and* surface degrees of freedom, and that the boundary symplectic structure is that of Chern-Simons theory. The phase space variables may be chosen to have the falloff required of asymptotically flat spacetimes at spatial infinity (the outer boundary). This completes the classical analogy with the true gravitational construction of Ref. [11]. (Note that the entire analogy may be constructed without reference to any covariant action: one may consider phase space variables and constraints on a spatial slice with inner and outer boundary, and impose any boundary conditions “by hand.”)

So far it appears that a difference from the gravitational case, apart from the identically vanishing Hamiltonian constraint [20], is that there is no Immirzi parameter γ ambiguity in the theory (5). This parameter plays an essential role [11] in fixing to $1/4$ the proportionality

constant between entropy and area. In fact, the Immirzi parameter *is* present in the canonical theory arising from (5). To see this, note that since the spatial dreibein e^{ai} is in general invertible (as for general relativity), we can construct the usual connection Γ_i^a via

$$\partial_{[a}e_{b]}^i = \epsilon^i_{jk}\Gamma_{[a}^j e_{b]}^k. \quad (6)$$

Now define the “scaled” canonically conjugate variables

$$\gamma A_a^i := \Gamma_a^i - \gamma K_a^i; \quad \gamma E_i^a := \frac{1}{\gamma} E_i^a, \quad (7)$$

where $K_a^i \equiv \Gamma_a^i - A_a^i$. This is exactly how the γ ambiguity enters in canonical gravity [15]. (Note however that the γ parameter can arise directly by introducing it into an action via a “ γ dual” of the four-dimensional spin connection [14]; in this sense, γ is more natural in general relativity than in the SU(2) theory (5), where this cannot be done.)

Now, since the entire theory (5) is the kinematical sector of general relativity, all the above quantization steps are *identical*. Therefore one can associate an entropy with the surface S which is proportional to its area A_S . Furthermore, γ may be fixed, as in [11], to get precisely $S = A/4$ in units of l_P .

This result seems a bit surprising, since as noted above there is no speed of light c in (5). However, as noted above, there are still two coupling constants λ and $\hbar$ in the quantum theory which may be combined to give a fundamental length.

The entropy may be interpreted similarly to the so called “entanglement entropy” calculated in any system, obtained by tracing over a portion of the available microscopic degrees of freedom [21]. Here as in [11], the bulk states are traced over to obtain an effective density matrix for the surface states. One important difference from other entanglement entropy calculations arises due to the quantum discreteness arising from the quantization: the “polymer” nature of the discrete geometry [22] naturally makes the entanglement entropy finite.

(II) In the spatial metric (q_{ab}) and extrinsic curvature (K_{ab}) variables for 3+1 gravity, a spatial two-surface S with spatial unit normal s^a is marginally outer trapped [23] (ie. is an apparent horizon) if

$$(q^{ab} - s^a s^b)(K_{ab} + D_a s_b) = 0, \quad (8)$$

where D_a is the covariant derivative associated with q_{ab} . This equation expresses the statement that the future outward expansion of light rays vanishes on S . In terms of the conjugate momentum $\tilde{\pi}^{ab} \equiv \sqrt{q}(K^{ab} - Kq^{ab})$, the equation may be succinctly written as

$$\tilde{\pi}^{ab} s_a s_b = \sqrt{q} D_a s^a. \quad (9)$$

This phase space condition has non-trivial solutions on generic spatial slices of black hole spacetimes. At late times, the radius of the apparent horizon determined by this equation coincides with the radius of the black hole event horizon. Therefore, it is the natural phase space condition to impose on an inner spatial boundary in the Hamiltonian theory, (if one wishes to follow the route of identifying black hole entropy as an entropy associated with horizon surface degrees of freedom, along the lines of [7] or [11]).

We now ask whether the condition (2) is the same as the apparent horizon equation (8). A straightforward argument shows that the answer is no. The main point to note is that the

apparent horizon equation (8) depends on the extrinsic curvature of S as embedded in a spatial surface Σ . On the other hand, eqn. (2) contains information only about the extrinsic curvature of Σ through A_a^i ; no derivatives of the spatial normal s^a of S appear. Indeed, information about s^a in (2) only appears in the projection $(q^{ab} - s^a s^b)$, and in the area 2-form $\underline{\sigma}_{ab}$. Now, because no information about the extrinsic curvature of S as embedded in Σ is present in (2), this phase space condition does not contain information about the expansion of light rays on the surface. Therefore it cannot be the apparent horizon equation. Thus, the fact that S is taken to be a trapped surface in [11] does not actually enter the quantum black hole entropy calculation.

(III) There is at least one other boundary condition, induced by adding a different surface term to the gravitational action, which also leads by arguments similar to the above, to an entropy proportional to area. Consider the action

$$S[e, A] = \frac{1}{8\pi G_N} \int_M \text{Tr} [e \wedge e \wedge F(A)] + \frac{1}{8\pi G_N} \frac{A_S}{4\pi} \int_{\partial M} \text{Tr} [e \wedge F(A)]. \quad (10)$$

This is the gravitational action (1), but with a “BF” theory surface term. The symplectic structure arising from this action also has a surface contribution, but it is now that of BF theory rather than Chern-Simons theory. Functional differentiability of the action now requires the condition

$$(\underline{D} \wedge e)_{ab} = -\frac{2\pi}{A_S} \underline{\sigma}_{ab}, \quad (11)$$

which fixes the connection on S in a different way than in (2).

The quantization procedure is unchanged, except that the new boundary condition (11) must be imposed as a quantum condition. This may be done in a form identical to (4), with the $\hat{\underline{e}}$ part acting on volume states and the Gauss law part $\underline{D} \wedge \hat{\underline{e}}$ acting on surface states:

$$\left(1 \otimes \frac{2\pi\gamma}{A_S} (\underline{D} \wedge \hat{\underline{e}})_{ab} \cdot r + \hat{\underline{e}}_{ab} \cdot r \otimes 1 \right) \Psi_V \otimes \Psi_S = 0 \quad (12)$$

The difference from the Chern-Simons case is that the surface Gauss law is now generated by $\underline{D} \wedge e_{ab}$ rather than $\underline{E}_{ab}$. What is unchanged is that the Gauss law has sources where edges of the volume spin network state puncture the surface.

The main question now is whether the entropy calculated by counting spin network states solving this new constraint, subject to their being eigenstates of area with eigenvalue within $\pm l_P^2$ of A_S , also gives an entropy proportional to the surface’s area. Let us first recall the two ingredients [11] in this calculation for the Chern-Simons boundary condition: (a) for a set of punctures $P = \{j_{p1}, \dots, j_{pn}\}$, the number of solutions N_P of (4), for a large number of punctures, is

$$N_P \sim \prod_{j_p \in P} (2j_p + 1), \quad (13)$$

and (b) the eigenvalue of the area operator for the set P is

$$A_P \sim l_P^2 \sum_{p \in P} \sqrt{j_p(j_p + 1)}. \quad (14)$$

The entropy is obtained by counting all sets P compatible with the area A_P . The ratio S/A is maximized for all j_p equal and of spin $1/2$. In this case $S/A = c \ln 2/\sqrt{3}$, where c is a constant.

For the boundary condition (11) to give the same result, the degeneracy (13) must be the same, since the area spectrum is obviously unchanged. We argue that this is the case: for a single edge of spin j piercing S , the associated degeneracy is at least the usual $(2j+1)$ from angular momentum considerations. Now, looking at the quantum constraints (4) and (12), it is apparent that this degeneracy has the same source, namely the action of $\hat{\mathcal{G}}_{ab}$ on volume states. Thus the degeneracy originates on the right hand side of the respective Chern-Simons or BF Gauss laws. This suggests that it *is* the same in the two conditions, because its source is the same. A proof of this conjecture, similar to that for the Chern-Simons case [24], may be possible.

Our observations suggest that the interesting framework for calculating black hole entropy developed in Ref. [11] is general enough to encompass actions and boundary conditions other than the specific ones considered there. Furthermore, since the condition that the two-boundary S is a marginally trapped does not enter the entropy calculation, the result can also hold for any boundary in Σ , even one that has trapped regions outside it.

It is possible to apply the general setup, with boundary conditions arising from an action, to other theories described by connections, whether or not they are generally covariant, or have local dynamics. Unlike the example in (II) above, a purely topological example is 4-dimensional BF theory with a boundary Chern-Simons term

$$S = \frac{1}{8\pi\lambda} \int_M \text{Tr}[B \wedge F(A)] + \frac{1}{8\pi\lambda} \frac{A_S}{4\pi} \int_{\partial M} \text{Tr} \left[A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right]. \quad (15)$$

Here B is a dimensionless $\text{SU}(2)$ valued two-form field, and the coupling constant λ has the same dimension as in (5). The same boundary condition (2) is induced by requiring a well defined variational principle. However the kinematics and dynamics are different. The bulk spatial diffeomorphism constraint is replaced by the flat connection condition, which renders the bulk part of the theory entirely topological [25,26]. Spin network states are again solutions to the Gauss law, but now the flat connection constraint “collapses” the spin networks everywhere except on non-trivial bulk topology, and on the boundary punctures where (2) holds. The Hilbert space divides up into surface and bulk states as before. The entropy associated with the surface states can again be calculated to give similar answers.

This outline for BF theory with a Chern-Simons boundary term lends further support to the statement that one can always ascribe an entropy to surfaces by counting the number of surface states. It would be interesting to produce surface boundary conditions, such that when quantized as in (4), the entropy turns out *not* to be proportional to area, whether or not the theory is generally covariant. (A large class of boundary conditions for the generally covariant case for theories with connection variables are considered in ref. [27].)

We close with some further comments: (i) Concerning quantization of area, if the action (1) is indeed the relevant one for the black hole sector of general relativity, then from a path integral point of view, quantization of the area A_S also follows from the standard argument for quantization of the Chern-Simons coupling, due to the transformation properties of the Chern-Simons term under large (Yang-Mills) gauge transformations. (ii) It would be interesting to study the general relativity action with the surface term that induces the

apparent horizon equation (8). Since this equation involves the connection rather than the curvature, the boundary term would not at first sight be gauge invariant. It may be possible to avoid this if eqn. (8) can be rewritten directly in terms of the curvature. (iii) It appears that kinematical considerations alone are not sufficient in themselves to justify the association of black hole entropy with surface entropy, since as we have seen, this may be done for generally covariant theories with no black hole solutions. (iv) An argument using the Immirzi parameter γ is essential for obtaining the correct proportionality constant $1/4$ in $S \sim A_S$ [11]. In the non-gravitational examples considered here, this parameter is also present in the two theories defined by (5) and (15), provided E_i^a is not degenerate. If so, the entropy associated with bounding surfaces can be fixed in the same way.

Acknowledgements This work was supported by the Natural Science and Engineering Research Council of Canada.

REFERENCES

- [1] A. Ashtekar, Phys. Rev. Lett. **57**, 2244 (1986); *Lectures on Non-perturbative Canonical Gravity*, (World Scientific, Singapore, 1991).
- [2] T. Jacobson and L. Smolin, Nucl. Phys. **B299** 295 (1988); V. Husain, Nucl. Phys. **B313**, 711 (1989).
- [3] C. Rovelli and L. Smolin, Nucl. Phys. **B331**, 80 (1990).
- [4] A. Ashtekar and C. J. Isham, Class. Quant. Grav. **9** 1433 (1992).
- [5] A. Ashtekar, J. Lewandowski, D. Marolf, J. Mourao, and T. Thiemann, J. Math. Phys. **36**, 6456 (1996), and references therein.
- [6] T. Thiemann, Phys. Lett. **B380**, 257 (1996).
- [7] S. Carlip, Phys. Rev. **D51**, 632 (1995); Class. Quant. Grav. **12**, 2853 (1995); See also A. Balachandran, L. Chandar, and A. Momen, Nucl. Phys. **B461**, 581 (1996).
- [8] L. Smolin, J. Math. Phys. **36**, 6417 (1995).
- [9] C. Rovelli Phys. Rev. Lett. **77**, 3288 (1996); Helv. Phys. Acta. **69**, 583 (1996).
- [10] K. Krasnov, Phys. Rev. **D55**, 3505 (1997); “On statistical mechanics of a Schwarzschild black hole”, Gen. Rel. Grav. (in press), gr-qc/9605047.
- [11] A. Ashtekar, J. Baez, A. Corichi and K. Krasnov, Phys. Rev. Lett. **80**, 904 (1998); A. Ashtekar and K. Krasnov, “Quantum geometry and black hole entropy,” gr-qc/9804039.
- [12] J. F. Plebanski, J. Math. Phys. **18**, 2511 (1977); J. Samuel, Pramana J. Phys. **28**, L429 (1987); T. Jacobson and L. Smolin, Phys. Lett. **B196**, 39 (1987).
- [13] F. Barbero, Phys. Rev. **D51**, 5498(1995); Phys. Rev. **D54**, 1492 (1996).
- [14] S. Holst, Phys. Rev. **D53**, 5966(1996).
- [15] G. Immirzi, “Quantum gravity and Regge calculus,” gr-qc/9701052; C. Rovelli and T. Thiemann, “The Immirzi parameter in quantum general relativity,” gr-qc/9605047.
- [16] C. Rovelli and L. Smolin, Phys. Rev. **D52**, 5743 (1995); J. Baez, Adv. Math. **117**, 253 (1996).
- [17] A. Ashtekar and J. Lewandowski, in *Knots and Quantum Gravity*, edited by J. Baez (Oxford U.P., Oxford, 1994); J. Baez, Lett. Math. Phys. **31**, 213 (1994).
- [18] C. Rovelli and L. Smolin, Nucl. Phys. **B442**, 593 (1995); S. Frittelli, L. Lehner, C. Rovelli, Class. Quant. Grav. **13**, 2921 (1996); K. Krasnov, “On the constant that fixes the area spectrum in canonical quantum gravity”, gr-qc/9709058.
- [19] A. Ashtekar, J. Lewandowski, Class. Quant. Grav. **14**, 55 (1997).
- [20] V. Husain and K. Kuchar, Phys. Rev. **D42**, 4070 (1990).
- [21] L. Bombelli, R. K. Kaul, J. Lee and R. D. Sorkin, Phys. Rev. D **34**, 373 (1987); M. Srednicki, Phys. Rev. Lett. **71**, 666 (1993).
- [22] A. Ashtekar, Int. J. Mod. Phys. **D5**, 629 (1996).
- [23] S. W. Hawking and G. F. R. Ellis, *The large scale structure of spacetime* (University of Cambridge Press, Cambridge, 1973).
- [24] R. K. Kaul and P. Majumdar, “Quantum black hole entropy,” gr-qc/9801080.
- [25] G. Horowitz, Commun. Math. Phys. **125**, 417 (1989).
- [26] V. Husain, Phys. Rev. **D43**, 1803 (1991).
- [27] V. Husain and S. Major, Nucl. Phys. **B500**, 381 (1997).
- [28] S. Carlip, “What we don’t know about BTZ black hole entropy,” hep-th/9806026.